

# On the Convergence Speed of Tetration

**Marco Ripà**

sPIqr Society, World Intelligence Network

Rome, Italy

e-mail: marcokrt1984@yahoo.it

**Published Online:** 15 Oct. 2018

**Abstract:** In 2011, in his book “La strana coda della serie  $n^n^{\dots^n}$ ”, M. Ripà analyzed some properties involving the rightmost figures of integer tetration, the iterated exponentiation  ${}^b a$ , characterized by an increasing number of stable digits for any base  $a > 1$ . A few conjectures arose about how many new stable digits are generated by any unitary increment of the hyperexponent  $b$ , and Ripà indicated this value as  $V(a)$  or “convergence speed” of  $a$ . In fact, when  $b$  is large enough,  $V(a)$  seems to not depend from  $b$ , taking on a (strictly positive) unique value, and many observations supported this claim. Moreover, we claim that  $V(a) = 1$  for any  $a \pmod{25} \in \{2, 3, 4, 6, 8, 9, 11, 12, 13, 14, 16, 17, 19, 21, 22, 23\}$  and  $V(a) \geq 2$  otherwise.

**Keywords:** Number theory, Power tower, Tetration, Chinese remainder theorem, Carmichael function, Euler’s totient function, Exponentiation, Integer sequence, Graham’s number, Convergence speed, Modular arithmetic, Stable digit, Rightmost digit.

**2010 Mathematics Subject Classification:** 11A07, 11F33.

## 1 Introduction

In the present paper, we introduce some conjectures involving the rightmost digits of the Tetration  ${}^b a = a^{a^{\dots}}$  ( $b$ -times) [1], observing that, when the hyperexponent  $b$  is sufficiently large and  $a \pmod{25} \notin \{0, 1, 5, 7, 10, 15, 18, 20, 24\}$ , the amount of new stable digits generated by any unitary increment of  $b$  is unitary as well: it depends only on the congruence modulo 25 of the base  $a$  [2].

This new result, if formally proved, would contribute to improve big numbers rightmost digits calculations, opening new scenarios in cryptography/cryptanalysis too [3-4].

## 2 Convergence Speed and Congruence (mod 25)

It is well known that, for any arbitrarily large  $n$ ,  ${}^b a$  originates a string of  $n$  stable figures, thus we can say that  ${}^b a$  is well-defined modulo  $10^n$ , for any  $b \geq b'(n, a)$  [2-5-6].

We can easily prove also that,  $\forall k \in \mathbb{N}$ ,  $a^{20 \cdot k+1} \equiv a \pmod{25}$ . In fact,  $\forall n \in \mathbb{N} \setminus \{0\}$ ,  $\lambda(n) \leq \varphi(n)$ . Thus,  $\lambda(25) = \varphi(25) = 20$ .

Let  $a$  be such that  $\gcd(a, 25) = 1 \Leftrightarrow \gcd(a, 5) = 1$ ,  $a^{\lambda(25)} \equiv 1 \pmod{25} \Rightarrow a^{20+1} \equiv a \pmod{25}$ . Hence  $a^{20 \cdot k+1} \equiv a \pmod{25}$ .

For any  $a$  such that  $a \equiv 5 \pmod{10}$ ,  $\forall m \in \mathbb{N} \setminus \{0, 1\}$ ,  $a^m \equiv 0 \pmod{25} \Rightarrow a^m \equiv a^{m+1} \pmod{25}$ . Therefore,  $a^2 \pmod{25} \equiv a^3 \pmod{25} \equiv \dots \equiv a^{20+1} \pmod{25} \equiv \dots \equiv a^{20 \cdot k+1} \pmod{25}$ .  $\square$

Let us now introduce the definition of “convergence speed” as it was originally presented by Ripà in his book about the rightmost digits of  ${}^b a$  [1].

**Definition 1:** Let  $a \in \mathbb{N} \setminus \{1\}$  be an arbitrary base which is not a multiple of 10 and let  $b \in \mathbb{N} \setminus \{0, 1\}$  be such that  ${}^{(b-1)} a \equiv {}^b a \pmod{10^d} \wedge {}^{(b-1)} a \not\equiv {}^b a \pmod{10^{(d+1)}}$ , where  $d \in \mathbb{N}$ , we consider  $V(a) \ni {}^b a \equiv {}^{(b+1)} a \pmod{10^{(d+V(a))}} \wedge {}^b a \not\equiv {}^{(b+1)} a \pmod{10^{(d+V(a)+1)}}$ .

For simplicity, from here on out, we refer to  $V(a)$  as the “convergence speed” of the natural base  $1 < a \not\equiv 0 \pmod{10}$  of the tetration  ${}^{(b \geq b')} a$ .

## 3 The Conjectures about $V(a)$

In this section we present the conjectures and a few remarks to point out their main implications.

**Conjecture 1:**  $\forall a \in \mathbb{N} \setminus \{1, 2\}$  such that  $a \not\equiv 0 \pmod{10}$ ,  $\exists b' < a \in \mathbb{N} \setminus \{0\} \ni \forall b \geq b'$ ,  $V(a) \in \mathbb{N} \setminus \{0\}$  is constant (see A317905 of the OEIS - ruling out the first term of the sequence [2]).

**Conjecture 2:** Assume  $b \in \mathbb{N} \setminus \{0, 1, 2\}$ ,  $\forall a \in \mathbb{N} \setminus \{1\}$  such that  $a \not\equiv 0 \pmod{10}$ ,  ${}^b a \equiv {}^{(b+1)} a \pmod{10^{((b-2) \cdot V(a))}}$ .

**Remark:** If Conjecture 2 holds, it follows that  $(b-2) \cdot V(a) \leq d + V(a)$ , hence  $\forall b > 3$ ,  $V(a) \leq \frac{d}{b-3}$  (e.g., if  $a = 143^{625}$  and  $b \geq 5$ ,  $4 = V(a) \leq \frac{0+6+6+5+\sum_{i=5}^b 4}{b-3} = \frac{17+(b-4) \cdot 4}{b-3}$  is true).

**Ripà's hypothesis:**  $\forall a \in \mathbb{N} \setminus \{1, 2\}$  such that  $a \not\equiv 0 \pmod{10}$ ,  $\exists b' < a \in \mathbb{N} \setminus \{0\} \ni \forall b \geq b'$ ,

$$\begin{cases} V(a) = 1 \Leftrightarrow a \pmod{25} \in \mathbb{C} = \{2, 3, 4, 6, 8, 9, 11, 12, 13, 14, 16, 17, 19, 21, 22, 23\} \\ V(a) \geq 2 \Leftrightarrow a \pmod{25} \in \mathbb{C} = \{0, 1, 5, 7, 15, 18, 24\} \end{cases}$$

**Remark:** It is very important to notice that, given  $a(\text{mod } 25) \in \mathbb{C}$  (or equivalently  $V(a) = 1$ ), it follows that  $V(a^m) \geq 2, \forall m = 5 \cdot n \in \mathbb{N} \setminus \{0\}$ , and  $V(a^m) = 1$  otherwise (for any  $m$  such that  $m(\text{mod } 10) \equiv \{1, 2, 3, 4, 6, 7, 8, 9\}$ ).

On the contrary, for any base such that  $a(\text{mod } 25) \in \mathbb{C}, V(a^n) \geq 2$ , since  $a^n(\text{mod } 25) \in \mathbb{C}$  too ( $\forall n \in \mathbb{N} \setminus \{0\}$ ). We point out that  $V(a) \geq 2 \Rightarrow a^{m+1}(\text{mod } 25) \equiv a(\text{mod } 25), \forall m = 4 \cdot n$ .

**Conjecture 3:**  $\forall v \in \mathbb{N} \setminus \{0\}, \exists a$ , not a multiple of 10, such that  $V(a) = v$ .

**Remark:** In order to prove Conjecture 3, it is sufficient to verify that, for any  $n$ -digits long base  $a := a_n \dots a_2 a_1$ , where  $a_1 = a_2 = \dots = a_n = 9, V(a = 9 \dots 9) = n (\forall b)$  (see [1], pp. 25-26).

From Ripà's hypothesis, it follows that  $a(n = 1) \in \mathbb{C} \Rightarrow V(a) = 1$  and  $a(n \geq 2) \in \mathbb{C} \Rightarrow V(a) \geq 2$ .

**Conjecture 4:** Let  $\text{len}(a(i))$  denote the length of the  $i$ -th term any (strictly positive) integer sequence  $a(n)$  constructed through the juxtaposition of integers,  $\forall i \in \mathbb{N}$  such that  $\text{len}(a(i)) \geq 2$ ,  $a^{(i)} a(i) \equiv a^{(i+1)} a(i+1) (\text{mod } 10^{\text{len}(a(i))})$ .

**Remark:** This conjecture was firstly introduced in 2011 [1] and two examples of this property are given by the sequences A317903 and A317824 of the OEIS [7-8].

## 4 Conclusion

It is not easy to provide a short proof of any of the conjectures introduced in Section 3 and this could be the subject of another paper that we hope to release in the near future.

We conclude with a very important question that we wish to answer:

“Is it possible to identify a new function

$\mathcal{R}(V(a)) := \min_R |\{V(a + k \cdot R) = V(a)\}|, \forall V(a) = n \in \mathbb{N} \setminus \{0\}$  and  $\forall k \in \mathbb{N}_0$

(e.g.,  $\mathcal{R}(V(a) = 1) = \mathcal{R}(1) = 25$  by Ripà's hypothesis)?”.

Any original contribute to help us to prove the aforementioned conjectures would be appreciated.

## References

- [1] Ripà, M., La strana coda della serie  $n^n \dots n^n$ . *UNI Service*, Trento, 2011.
- [2] Sloane, N. J. A., *The Online Encyclopedia of Integer Sequences*, Web. 10 Aug. 2018, <http://oeis.org/A317905>
- [3] Germain, J., On the Equation  $a^x \equiv x (\text{mod } b)$ . *Integers: Learning, Memory, and Cognition*, 9(6), 2009, 629–638.
- [4] Urroz, J. J., Yebra J. L. A., On the Equation  $a^x \equiv x (\text{mod } b^n)$ , *Journal of Integer Sequences*, 8(8), 2009.

- [5] User26486 & Milo Brandt, *Mathematics Stack Exchange*, Do the last digits of exponential towers really converge to a fixed sequence?, Web. 22 Feb. 2015, <https://math.stackexchange.com/questions/1159995/do-the-last-digits-of-exponential-towers-really-converge-to-a-fixed-sequence>
- [6] Vladimir & Doctor Jacques, *The Math Forum*, Modular Exponentiation, Web. 26 Mar. 2005, <http://mathforum.org/library/drmath/view/51625.html>
- [7] Sloane, N. J. A., *The Online Encyclopedia of Integer Sequences*, Web. 10 Aug. 2018, <http://oeis.org/A317903>
- [8] Sloane, N. J. A., *The Online Encyclopedia of Integer Sequences*, Web. 10 Aug. 2018, <http://oeis.org/A317824>